arXiv:1201.6683v1 [math.AP] 31 Jan 2012

HOMOGENIZATION OF THE BOUNDARY VALUE FOR THE DIRICHLET PROBLEM

KI-AHM LEE AND HENRIK SHAHGHOLIAN

Abstract. The Dirichlet problem with oscillating boundary data is the subject of study in this paper. It turns out that due to integral representation of such problems we can reduce the study to the case of surface integrals of rapidly oscillating functions, and their limit behavior:

$$\lim_{\varepsilon\to 0}\int_\Gamma g\left(y,\frac{y}{\varepsilon}\right)d\sigma_y,$$

where $g(x,y)$, represents the boundary value in the Dirichlet problem. The lower dimensional character of the surface Γ produces unexpected and surprising effective limits. In general, the limit of the integral depends strongly on the sequence $\varepsilon=\varepsilon_j$ chosen. Notwithstanding this, when the surface does not have flat portions with *rational directions* a full averaging takes place and we obtain a unique effective limit in the above integral.

The results here are connected to recent works of D. Gérard-Varet and N. Masmoudi, where they study this problem in combination with homogenization of the operator, in convex domains.

Contents

1. Introduction 2
1.1. Background 2
1.2. Heuristics 3
1.3. Plan of the paper 5
2. Definitions and Notation 5
2.1. Averaging and Ergodic theory 5
3. Technical preliminaries 7
4. Surface integrals of oscillating functions 11
5. The case of Layered densities, almost periodicity, and ergodicity 13
5.1. Layered Materials 13
5.2. Almost Periodic Case 13

H. Shahgholian has been supported in part by Swedish Research Council. K. Lee has been supported by Korea-Sweden Research Cooperation Program. This project is part of an STINT (Sweden)-NRF (Korea) research cooperation program.

5.3. Ergodic Case 13
6. Applications to Partial Differential Equations 13
6.1. Dirichlet problem: Elliptic Case 13
6.2. Dirichlet problem: Parabolic Case 15
6.3. Neumann problem 15
7. Examples and illustrations 17
7.1. Example 1 18
7.2. Example 2 20
References 21

1. INTRODUCTION

1.1. Background.

In this paper we consider Dirichlet and related problems, with rapidly oscillating data. Although we treat the case of Laplace equation, our analysis straightforwardly extends to general cases for equations that admit a Poisson/Green representation. Such integral representation, in turn, reduces the study of the problem to the corresponding integral equation, and hence the analysis of the integral of rapidly oscillating functions becomes central.

To fix ideas, let Γ be a C^1 surface in $\mathbb{R}^n$ ($n \geq 2$), not necessarily bounded. We assume $g(x,y)$ is integrable in both variables over Γ, and 1-periodic in y-variable i.e., $g(x, y+k) = g(x,y)$ for $k \in \mathbb{Z}^n$. In this paper, we shall study possible limit behaviors of the integral

$$\lim_{\varepsilon\to 0} \int_\Gamma g\left(y, \frac{y}{\varepsilon}\right) d\sigma_y, \tag{1}$$

and we shall prove that under mild conditions on the surface Γ there is an effective limit as ε tends to zero.

In general we show that the above integrals stay within the bounds of the interval $\left[\int_\Gamma \overline{g}_*(y,\nu_y), \int_\Gamma \overline{g}^*(y,\nu_y)\right]$ where $\overline{g}_*, \overline{g}^*$ are defined below, as the infimum respectively supremum of the average-integrals of $g(y,\cdot)$ over closed loops of the plane $\{x : x\cdot\nu_y = 0\}$ on the torus; here ν_y denotes the normal vector of Γ at y.

To illuminate the application of this to the Dirichlet problem, let us consider a bounded C^1-domain $\Omega \subset \mathbb{R}^n$ ($n \geq 2$). Let further $g(x,y)$ and $f(x,y)$ be continuous in both variables, and periodic in y. Let u_ε be a solution to the Dirichlet problem

$$(P_\varepsilon) \qquad \triangle u_\varepsilon = -\mu_\varepsilon \quad \text{in } \Omega, \qquad u_\epsilon = g(x, x/\varepsilon) \quad \text{on } \partial\Omega,$$

where $\mu_\varepsilon = f(x, x/\varepsilon)\chi_{\Gamma_0} d\sigma_x$, and Γ_0 is a C^1 surface compactly in Ω.

Since solutions to this problem can be represented by surface integrals of g and f (through Poisson and Green functions), we may directly apply our results. Hence, a by-product of our surface integral homogenization, is nontrivial limit scenarios, as ε tends to zero, of equation (P_ε) and its solution.

It is noteworthy that, through integral equations of Fredholm type, or just standard functional minimization, our technique applies to homogenizations with oscillating Neumann data (see Section 6.3). For fully nonlinear operator the Neumann problem has been treated in [CKL], see also [BDLS].

Another issue that occurs along such analysis is the study of the speed of convergence. In problems, where governing partial differential equations have oscillating coefficients, one uses the standard method of expansion

$$u_\epsilon(x) = u_0 + \epsilon u_1(x, x/\epsilon) + \epsilon u_{bl,\epsilon} + O(\epsilon),$$

in an appropriate space. Here $u_{bl,\epsilon}$ is the so-called boundary layer term, and finding it is part of the problem. The function $u_{bl,\epsilon}$ will then solve a Dirichlet problem with oscillating boundary data. We refer to [AA] for some backgrounds and to [GM] for recent developments in the topic.

Remark 1.1. *It is noteworthy that our approach applies to general equations of the type*

$$div(A(x/\varepsilon)\nabla u) = f(x/\varepsilon)$$

with oscillating Dirichlet/Neumann data. This, even though straightforward, becomes quite technical and is therefore outside the scope of this paper. Thus for clarity of the exposition we shall treat the Laplacian case, only.

1.2. **Heuristics.** Let us set $x = y (\text{mod } 1)$ if $x \in Q_1^+ = \{0 \leq x_i < 1\}$, and $y - x \in \mathbb{Z}^n$, and adopt the standard notation from homogenization, as well as that of ergodic theory.

For clarity of the ideas, let us also deal with problem (P_ε) in the case $f \equiv 0$, and see how it reduces to the integral averaging, and then how the limit integral is obtained.

Since harmonic functions can be represented using Poisson kernel $P(x, y)$,

$$u_\epsilon(x) = \int_{\partial\Omega} P(x, y) g(y, y/\epsilon)\, d\sigma_y,$$

we must then analyze the behavior of this integrals as ε tends to zero.

For $x \notin \partial\Omega$, $P(x,y)$ is continuous and we can rewrite the integral

$$u_\epsilon(x) \approx \sum_j P(x,y^j)|\partial\Omega \cap Q_{r_j}(y^j)| \fint_{\partial\Omega\cap Q_{r_j}(y^j)} g(y^j, y/\epsilon)\, d\sigma_y, \tag{2}$$

where r_j is small enough, independent of ε, and $Q_{r_j}(y^j)$ is a cube with size r_j and center y^j. This obviously brings us to integral (1). Observe that at this stage we cannot replace the second variable of g with y^j, due to rapid oscillation for $\epsilon \approx 0$.

Further assume that $\partial\Omega \cap Q_{r_j}(y^j)$ is so flat that at ϵ-scale it is approximately ϵ^2-away from its tangent plane; this requires a C^2-graph locally, but our techniques/proofs work for C^1-surfaces.

The idea now is to cover the boundary $\partial\Omega \cap Q_{r_j}(y^j)$ with finite many, and small enough cubes, so that each part of the surfaces is as flat as we want to. In particular we have

$$\fint_{\partial\Omega\cap Q_{r_j}(y^j)} g(y^j, y/\epsilon)\, d\sigma_y = \sum_i \fint_{\partial\Omega\cap Q_{\varepsilon_i r_j}(y^{i,j})} g(y^j, y/\epsilon)\, d\sigma_y,$$

where ε_i is to be chosen small enough. This part is slightly more delicate, and needs extra care. We would also prefer to take $\varepsilon_i = \sqrt{\varepsilon}$. The trouble is that this might not work, as the points are changing, and we may loose lots of information about the behavior of the surface $\partial\Omega \cap Q_{\varepsilon_i r_j}(y^{i,j})$. Indeed, an scaling of the integral gives

$$\fint_{\partial\Omega\cap Q_{(\varepsilon_i/\varepsilon) r_j}(y^{i,j})} g(y^j, y)\, d\sigma_y,$$

and the question is whether this integral will converge to the mean $\fint_{Q_1^+} g(y^j, y) dy$. This would happen exactly when the normal vector of the surface $\partial\Omega$ at $y^{i,j}$ is irrational. In other words the surface $\partial\Omega \cap Q_{(\varepsilon_i/\varepsilon) r_j}(y^{i,j})$ (which is almost a plane) will foliate the n-dimensional torus, as ε tends to zero, provided we have chosen $\varepsilon_i/\varepsilon \to \infty$, and the normal vector of the surface $\partial\Omega$ at $y^{i,j}$ is irrational. In particular

$$\lim_{\varepsilon\to 0} \fint_{\partial\Omega\cap Q_{(\varepsilon_i/\varepsilon) r_j}(y^{i,j})} g(y^j, y)\, d\sigma_y = \overline{g}(y^j). \tag{3}$$

If the set of points on $\partial\Omega$ with rational direction has zero surface measure (this may fail if there are so-called flat spots with rational directions) then we could cover the boundary with small cubes centered at points with irrational normals. From here combining (2)-(3)

we obtain

$$\lim_{\varepsilon\to 0} u_\varepsilon(x) = \int_{\partial\Omega} P(x,y)\overline{g}(y)\, d\sigma_y .$$

It is apparent that if there is a flat portion of the boundary with rational normal, then a full foliation cannot take place at such portions. Consequently, the resulting (mod 1) surface will be a close simple curve over the n-dimensional torus. Hence different sequence of ϵ may give different shifts of this close loop, and hence the possibility of a parameter family of values, in between $[\overline{g}_*(y,\nu_y), \overline{g}^*(y,\nu_y)]$.

Remark 1.2. ***A word of caution:*** *As ε tends to zero, one may obviously rescale the integral, by a change of variables, as we did in our heuristic explanations above. This scaling makes the surface to be scaled (we assume the origin is NOT on the surface) so as to disappear in the limit. This naturally would make it impossible to compute the limit integral. However, the periodicity of the function g in its second variable, implies that we can bring back the surface so as it passes through the fundamental cube Q_1^+ using (mod1) argument. In particular this means that despite the integration is on a fixed surface, the surface itself will start jumping forth and back due to the variable ε.*

It would be a good idea for the reader to consider simple examples such as integration over a line segment in the plane, by varying the normal direction of the plan from rational to irrational.

1.3. **Plan of the paper.** In the next section we shall introduce all definitions and notations. We take care of some technicalities in Section 3. We shall formulate our main results concerning surface integration in Section 4, and its generalization to other type of functions, such as layered-densities, almost periodic functions will appear in Section 5. Several interesting applications to PDE are mentioned in Section 6, and several Examples are also given in Section 7.

2. DEFINITIONS AND NOTATION

2.1. Averaging and Ergodic theory.

Definition 2.1. *Let ν be a vector in $\mathbb{R}^n$, z a fixed given point, and $Q_r(z) = \{x : |x_i - z_i| < r\}$. Let $g(x,y)$, be integrable in both variables over the plain $\{(y-z)\cdot\nu = 0\}$, and periodic in y-variable. We define*

$$\overline{g}^*(z,\nu) := \overline{\lim_{\varepsilon\to 0}}\,\overline{\lim_{r\to\infty}} \fint_{\{(y-z)\cdot\nu=0\}\cap Q_r(z)} g\left(z,\frac{y}{\varepsilon}\right) d\sigma_y ,$$

and

$$\overline{g}_*(z,\nu) := \underline{\lim_{\varepsilon\to 0}}\,\underline{\lim_{r\to\infty}} \fint_{\{(y-z)\cdot\nu=0\}\cap Q_r(z)} g\left(z,\frac{y}{\varepsilon}\right) d\sigma_y .$$

Later we shall consider cases where $\nu = \nu_z$, *is the normal vector at* z *on a given surface* Γ. *We also define the average of* g *as*

$$\overline{g}(z) = \fint_{Q_1^+} g\,(z, y)\ dy,$$

for $Q_1^+ = \{x : 0 \leq x_i \leq 1\}$.

Henceforth we shall assume all vectors have length one, unless otherwise stated. We shall also without loss of generality assume that the surface is orientable, and fix a consistent choice of normal in clockwise direction (this choice is obvious if Γ is the boundary of a domain).

It should be noted that if the set $\{(y - z) \cdot \nu = 0\}(mod\ 1)$ foliates the cube Q_1^+ then the integral

$$\lim_{r\to\infty} \fint_{\{(y-z)\cdot\nu=0\}\cap Q_r(z)} g\left(z, \frac{y}{\varepsilon}\right) d\sigma_y$$

converges to the average $\overline{g}(z)$, and this happens exactly when ν is irrational direction (see Lemma 3.2). When the set $\{(y - z) \cdot \nu = 0\}(mod\ 1)$ does not foliate the cube Q_1^+ (or the n-dimensional torus) then we shall get the limit as an integral over a closed loop flow on the torus. In particular the value of the integral exists, and depends on ν and ϵ. For different values of ϵ this loop translates over the torus and will give rise to supremum respectively infimum values of the integrals, as defined above by $\overline{g}^*, \overline{g}_*$, respectively.

Remark 2.1. *It should be remarked that in the definitions of the above averages* $\overline{g}^*, \overline{g}_*$ *we could have replaced*

$$\fint_{\{(y-z)\cdot\nu=0\}\cap Q_r(z)} g\left(z, \frac{y}{\varepsilon}\right) d\sigma_y,$$

by

$$\fint_{\{(y-z)\cdot\nu=0\}\cap D} g\left(z, \frac{y}{\varepsilon}\right) d\sigma_y,$$

for any domain D *containing the point* z. *The reason for this is that the set* $T = \{(y - z) \cdot \nu = 0\}(mod\ 1)$ *either foliates the whole* n*-dimensional torus, or it is a closed simple curve on the torus. In either case, when* ϵ *tends to zero, the piece of plane* $\{(y - z) \cdot \nu = 0\} \cap D$ *will have the same effect as* T.

Since the direction of the normal of the plane $\{(y - z) \cdot \nu = 0\}$ will play a crucial role in our analysis, we introduce proper definitions well-known in Ergodic theory. For readers' convenience we also prove some of these well-known results here.

Definition 2.2. *(IDDC)*

1. *The vector $\nu \in \mathbb{S}^{n-1}$ is a rational direction if there is $m \in \mathbb{Z}^n$ such that $\nu = \frac{m}{|m|}$.*
2. *The vector $\nu \in \mathbb{S}^{n-1}$ is an irrational direction if ν is not a rational direction.*
3. *A smooth surface Γ satisfies Irrational Direction Dense Condition (IDDC) if*

$$\sigma_\Gamma\left(\Gamma \setminus \{x \in \Gamma : \nu_x \text{ is a irrational direction. }\}\right) = 0$$

for the surface measure σ_Γ. Here ν_x denotes the normal to Γ at x.

3. Technical preliminaries

In this section we shall recall some standard facts from Ergodic theory, and also state and prove some averaging results that will be needed for the proof of the main theorems.

The first lemma is a version of Weyl's Lemma and is well-known fact about uniform distribution.

Lemma 3.1 (Weyl's Lemma)**.** *Let $\nu = (\nu_1, \cdots, \nu_d) \in \mathbb{R}^d$ $(d \geq 1)$, be such that $(1, \nu_1, \cdots, \nu_d)$ is an irrational direction. Then, for $h \in L^2(0,1)$, and $a_k = \sum_{i=1}^d k_i \nu_i (mod\ 1)$ we have*

$$\frac{1}{|I_N|}\sum_{k\in I_N} h(a_k) \to \int_0^1 h(t)dt, \qquad k = (k_1, \cdots, k_d)$$

as $N \to \infty$. Here $I_N = \{k \in \mathbb{Z}^d : |k_i| \leq N, i = 1, \cdots, d\}$.

That $(1, \nu_1, \cdots, \nu_d)$ is an irrational direction, means that at least one of the numbers ν_j is irrational. In this paper $d = n-1$ is the only case that is used.

Proof. Set

$$\mu_{\nu,N} = \frac{1}{|I_N|}\sum_{k\in I_N} \delta_{k\cdot\nu}$$

to be the sum of Dirac delta functions. First, let us consider the case $h = h_m = e^{2\mathrm{i}\pi m t}$ and $m \neq 0$. By the assumption there is ν_l, which is an irrational number, and hence

$$(4) \qquad \left|\mu_{\nu,N}(h_m)\right| = \frac{1}{|I_N|}\Pi_{j=1}^d \left|\sum_{k_j=-N}^{N} e^{2\mathrm{i}\pi m k_j \nu_j}\right| \leq \frac{C}{2N\left|1 - e^{2\mathrm{i}\pi m \nu_l}\right|} \quad \to_{N\to\infty} 0,$$

since $m\nu_l$ cannot be an integer i.e. $1 - e^{2\mathrm{i}\pi m\nu_l} \neq 0$. Now by Fourier expansion (extending h as 1-periodic function), for some $b_m \in \mathbb{R}$,

$$h(s) = \int_0^1 h(t)dt + \sum_{m=1} b_m e^{2\mathrm{i}\pi ms},$$

which in combination with (4) results in

$$\mu_{\nu,N}(h) = \int_0^1 h(t)dt + \sum_{m=1}^{\infty} b_m \mu_{\nu,N}(e^{\mathrm{i}2\pi ms}) \to \int_0^1 h(t)dt.$$

□

Lemma 3.2. *Let g, $\overline{g}^*$, $\overline{g}_*$, and $\overline{g}$ be as in Definition 2.1. Then the following hold:*

(i) If ν_x is an irrational direction, then

$$\overline{g}^*(x,\nu_x) = \overline{g}_*(x,\nu_x) = \overline{g}(x).$$

(ii) The following inequalities always hold

$$\overline{g}^*(x,\nu_x) \geq \overline{g}(x) \geq \overline{g}_*(x,\nu_x).$$

Proof. First note that we can replace the integral in the definition of $\overline{g}_*$, and $\overline{g}^*$ by

$$\fint_{\{(y-x)\cdot\nu=0\}\cap Q_N} g\left(x, \frac{y}{\varepsilon}\right) d\sigma_y,$$

with N positive integers, and $N \to \infty$. This simplifies the matter slightly.

Set $\Pi = \Pi(\nu, x) = \{(y-x)\cdot\nu = 0\}$. We may without loss of generality assume $\nu = (1, \nu')$. Since ν is irrational direction, $\Pi(mod\ 1)$ will foliate the unit cell, according to standard foliation theory and results in integral average over Q_1^+. To see this we assume (by using periodicity) that $x \in Q_1^+$ and that the plane Π cuts the x_1-axis (or some other axis). let a_0 be the point of intersection between the x_1-axis and this plane, so that $\Pi(\nu, x) = \{x : x_1 = a_0 - \nu' \cdot x'\}$, and $(1, \nu')$ is irrational (by the assumption), and $0 \leq a_0 < 1$ due to periodicity of $g(x, y)$ in y. Having fixed ν, we shall now use the notation $\Pi_t = \Pi(\nu, te_1)$, the plane with normal ν through the point $(t, 0')$.

For $k' \in \mathbb{Z}^{n-1}$ define $S_{k'}$ to be the cylinder generated by the x_1-axis and the $(n-1)$-dimensional cube $k' + \{y : 0 < y_i < 1,\ i = 2, \cdots, n\}$ (these are cubes in $\mathbb{R}^{n-1}$ translated by the vector k'). Set further $a_{k'} = a_0 - k' \cdot \nu' (mod\ 1)$ $(k' = (k_2, \cdots, k_n))$, then $\tilde{\Pi}_{a_{k'}} = (\Pi \cap S_{k'})\ (mod\ 1)$.

Let $N > 0$ be a large number, and $I'_N = \{k' \in \mathbb{Z}^{n-1} : |k_i| \leq N\}$. Then by periodicity of $g(x, \cdot)$

$$\int_{Q_N(x)\cap\Pi} g\,(x, y)\,d\sigma_y = \sum_{k'\in I'_N} \int_{\tilde{\Pi}_{a_{k'}}} g\,(x, y)\,d\sigma_y.$$

Set now $w(a_{k'}) = w(x, a_{k'}) := \int_{\tilde{\Pi}_{a_{k'}}} g\,(x, y)\,d\sigma_y$. Then

$$\fint_{Q_N\cap\Pi} g\,(x, y)\,d\sigma_y = \frac{1}{|\nu|}\frac{1}{|I'_N|} \sum_{k'\in I'_N} w(a_{k'}),$$

where we have used $|Q_N \cap \Pi| = (2N)^{n-1}|\nu| = |I'_N||\nu|$.

Since ν is irrational, from Lemma 3.1 we conclude

(5)
$$\lim_{N\to\infty} \fint_{Q_N\cap\Pi} g\,(x, y)\,d\sigma_y = \frac{1}{|\nu|}\int_0^1 w(t)dt = \frac{1}{|\nu|}\int_0^1 \int_{\tilde{\Pi}_t} g(x, y)d\sigma_y dt = \overline{g}(x),$$

where $\tilde{\Pi}_t = \Pi_t \cap S_{0'}$ (*mod* 1). Therefore $\lim_{r\to\infty} \fint_{\{(y-x)\cdot\nu=0\}\cap Q_r} g\left(x, \frac{y}{\varepsilon}\right) d\sigma_y = \overline{g}(x)$ which is independent of ε. Now we have conclusion (i) from the definition of $\overline{g}^*(x, \nu)$ and $\overline{g}_*(x, \nu)$.

To prove (ii) let us suppose that ν is a rational direction, otherwise the conclusion follows by the equality in (i). Since ν is rational, it is not hard to see that the restriction of $g(x, y)$ on this hyperplane will be periodic with period T, say; for $g(x, y/\varepsilon)$ the period will be εT. In particular the integral can be seen as integration over a spiral-like plane on the torus (a closed loop). The two dimensional case can be illustrated by a curve on the torus that loops over itself. In particular the limit integral (w.r.t. r) in the definition of $\overline{g}_*, \overline{g}^*$ can be replaced by

$$\overline{g}_*(x, \nu_x) = \underline{\lim}_{\varepsilon} \fint_{\{(y-x)\cdot\nu=0\}\cap Q_{\varepsilon T}} g\left(x, \frac{y}{\varepsilon}\right) d\sigma_y = \underline{\lim}_{\varepsilon} \fint_{\{(y-x_\varepsilon)\cdot\nu=0\}\cap Q_T} g\,(x, y)\,d\sigma_y,$$

$$\overline{g}^*(x, \nu_x) = \overline{\lim_{\varepsilon}} \fint_{\{(y-x)\cdot\nu=0\}\cap Q_{\varepsilon T}} g\left(x, \frac{y}{\varepsilon}\right) d\sigma_y = \overline{\lim_{\varepsilon}} \fint_{\{(y-x_\varepsilon)\cdot\nu=0\}\cap Q_T} g\,(x, y)\,d\sigma_y.$$

An observation here is that the supremum value of the mean integral w.r.t. ϵ actually is taken for some value ϵ_0, and naturally for many other values, due to periodicity in ϵ. One can think of the situation as parallel planes in $\mathbb{R}^n$ with fixed distance ϵ_1 from each other are moving, simultaneously by keeping a fixed distance between them, in the orthogonal direction of the plane (*mod* 1), when ϵ changes. Obviously, these planes foliate Q_1^+ once ϵ ranges $[0, 1)$.

From here one deduces

$$\fint_{Q_1^+} g(x,y)dy = \fint_0^{\epsilon_1} d\epsilon \fint_{\Pi(\nu,x_\epsilon)(mod\ 1)} g(x,y)d\sigma_y$$

$$\leq \fint_{\Pi(\nu,x_{\epsilon_0})(mod\ 1)} g(x,y)d\sigma_y \leq \overline{g}^*(x,\nu).$$

Similarly

$$\fint_{Q_1^+} g(x,y)dy \geq \overline{g}_*(x,\nu).$$

□

In our analysis of the limit behavior of the integral (1) we will use Definition 2.1 in a slightly different way. Next lemma will give us a hint in that direction.

Lemma 3.3. *Let g be as before, $\Pi(\nu,z) = \{(y-z)\cdot\nu = 0\}$, and $z \in \Gamma$. Then for any $R_\varepsilon \nearrow \infty$ (as $\varepsilon \searrow 0$) we have*

$$\limsup_{\varepsilon\to 0} \fint_{\Pi(\nu,z)\cap Q_{\varepsilon R_\varepsilon}(z)} g(z,y/\varepsilon)d\sigma_y \leq \overline{g}^*(z,\nu),$$

and

$$\liminf_{\varepsilon\to 0} \fint_{\Pi(\nu,z)\cap Q_{\varepsilon R_\varepsilon}(z)} g(z,y/\varepsilon)d\sigma_y \geq \overline{g}_*(z,\nu).$$

In case ν is irrational we obtain

$$\lim_{\varepsilon\to 0} \fint_{\Pi(\nu,z)\cap Q_{\varepsilon R_\varepsilon}(z)} g(z,y/\varepsilon)d\sigma_y = \overline{g}(z).$$

We leave the the reader to verify this simple fact.

We shall next make the previous lemma even more general by letting the plane be replaced by very smooth surfaces. Let Γ be a smooth surface, with module of continuity $\tau = \tau_\Gamma$ for its C^1 norm. Define

$$M_\epsilon = \sqrt{\min(\frac{1}{\tau(\varepsilon)}, \frac{1}{\varepsilon})} \nearrow \infty, \qquad \text{as } \epsilon \searrow 0. \tag{6}$$

Then

$$M_\epsilon \tau(\epsilon M_\epsilon) \leq \sqrt{\tau(\sqrt{\epsilon})} \to 0, \qquad \text{as } \epsilon \to 0. \tag{7}$$

Lemma 3.4. *Let g be as before, and Γ a smooth C^1 surface, with module of continuity $\tau = \tau_\Gamma$ for its C^1 norm. Let further $\rho_\varepsilon = \epsilon M_\epsilon$ where M_ϵ is as in*

(6). Then, for $z \in \Gamma$, and $\eta > 0$ there exists $\varepsilon_{\nu_z,\eta}$ such that for all $\varepsilon \leq \varepsilon_{\nu_z,\eta}$ we have

$$\overline{g}_*(z,\nu_z) - \eta \leq \fint_{\Gamma\cap Q_{\rho_\epsilon}(z)} g\left(z, \frac{y}{\varepsilon}\right) d\sigma_y \leq \overline{g}^*(z,\nu_z) + \eta.$$

Proof. Set $z_\varepsilon = \frac{z}{\varepsilon}\,(mod\ 1)$, and $\Gamma_z^\epsilon := \{x : \epsilon(x+z) \in \Gamma\}$. Then we have

$$\fint_{\Gamma\cap Q_{\rho_\epsilon}(z)} g\left(z, \frac{y}{\varepsilon}\right) d\sigma_y = \fint_{\Gamma^\varepsilon \cap Q_{M_\epsilon}(0)} g\left(z, z_\varepsilon + y\right) d\sigma_y,$$

where

$$\Gamma_z^\varepsilon \cap Q_{M_\varepsilon}(0) \subset \left\{ x \in Q_{M_\varepsilon}(0) : |x \cdot \nu_z| < \frac{\rho_\epsilon \tau(\rho_\varepsilon)}{\varepsilon} = M_\varepsilon \tau(\varepsilon M_\varepsilon) \right\}.$$

If we set $\Pi_{\nu_z} = \{x :\ x \cdot \nu_z = 0\}$, then by continuity of $g(z,.)$, and that for ε small enough the surface Γ_z^ε is $M_\varepsilon \tau(\varepsilon M_\varepsilon)(\approx 0)$ close to Π_{ν_z} in $B_{M_\varepsilon}(0)$ we will have

$$\fint_{\Gamma_z^\varepsilon \cap Q_{M_\epsilon}(0)} g\left(z, z_\varepsilon + y\right) d\sigma_y = \fint_{\Pi_{\nu_z} \cap Q_{M_\epsilon}(0)} g\left(z, z_\varepsilon + y\right) d\sigma_y + I,$$

where

$$I = \fint_{\Gamma_z^\varepsilon \cap Q_{M_\epsilon}(0)} \left(g\left(z, z_\varepsilon + \tilde{y}\right) - g\left(z, z_\varepsilon + y\right)\right) d\sigma_y.$$

Here $\tilde{y} = y + \nu_z |y| s_\varepsilon$ with $s_\varepsilon = \tau(\varepsilon M_\varepsilon)$. Estimating I we obtain $|I| \leq C\tau_g(|y|s_\varepsilon) \leq C\tau_g(M_\varepsilon s_\varepsilon) < \eta/2$, for ε small enough. Here τ_g is the module of continuity for g From here, and Lemma 3.3 the statements in the lemma follows, provided we have taken ε small enough depending on ν_z. □

4. Surface integrals of oscillating functions

Theorem 4.1. *Let Γ be a C^1 surface in $\mathbb{R}^n$, and $g(x,y)$ be integrable in x-variable over Γ, and continuos and 1-periodic in y in $\mathbb{R}^n$. Then*

$$\limsup_{\varepsilon\to 0} \int_\Gamma g\left(y, \frac{y}{\varepsilon}\right) d\sigma_y \leq \int_\Gamma \overline{g}^*(y,\nu_y) d\sigma_y,$$

and

$$\liminf_{\varepsilon\to 0} \int_\Gamma g\left(y, \frac{y}{\varepsilon}\right) d\sigma_y \geq \int_\Gamma \overline{g}_*(y,\nu_y) d\sigma_y.$$

Moreover if Γ satisfies IDDC then an effective limit exists and we have

$$\lim_{\varepsilon\to 0} \int_\Gamma g\left(y, \frac{y}{\varepsilon}\right) d\sigma_y = \int_\Gamma \overline{g}(y) d\sigma_y.$$

Proof. We shall prove the limit superior estimate only. The limit inferior estimates follows in a similar way. The last statement follows in an obvious manner.

Let us fix a small positive constant $\eta > 0$, to be decided later. Without loss of generality, we may assume Γ is bounded, and $g(x, y)$ is uniformly continuous function on $\Gamma \times \mathbb{R}^n$.

Next from Lemma 3.4, for each $z \in \Gamma$ we may take $\varepsilon_{\nu_z,\eta}$ such that the estimate

$$\overline{g}_*(z, \nu_z) - \eta \leq ⨍_{\Gamma \cap Q_{\rho_\varepsilon}(z)} g\left(z, \frac{y}{\varepsilon}\right) d\sigma_y \leq \overline{g}^*(z, \nu_z) + \eta. \tag{8}$$

holds for all $\varepsilon < \varepsilon_{\nu_z,\eta}$. Next we cover Γ by cubes $Q_{\rho_\varepsilon}(z)$ for all $z \in \Gamma$ and all $\rho_\varepsilon = \varepsilon M_\varepsilon$ with $\varepsilon < \varepsilon_{\nu_z,\eta}$. This is a fine cover of Γ, and therefore there is a *finite* disjoint sub-cover such that $\Gamma \setminus \cup_{j=1}^N Q_{\rho_{\varepsilon_j}}(z^j)$ has surface measure less than η. Here $z^j \in \Gamma$, and $\varepsilon_j = \varepsilon_{\nu_{z^j}}$. Next

$$\int_\Gamma g(y, y/\varepsilon) d\sigma_y \leq \sum_{j=1}^N \int_{\Gamma \cap Q_{\rho_{\varepsilon_j}}(z^j)} g(y, y/\varepsilon) d\sigma_y + \eta \|g\|_\infty,$$

and by (8) we obtain

$$\int_\Gamma g(y, y/\varepsilon) d\sigma_y \leq \sum_{j=1}^N \left(\overline{g}^*(z^j, \nu_{z^j}) + \eta\right) |\Gamma \cap Q_{\rho_{\varepsilon_j}}(z^j)| + \eta \|g\|_\infty$$

$$\leq \int_\Gamma \overline{g}^*(y, \nu_y) d\sigma_y + (1 + \|g\|_\infty + |\Gamma|)\eta < \int_\Gamma \overline{g}^*(y, \nu_y) d\sigma_y + \eta_0,$$

for any η_0 as small as we wish, provided η is small enough.

In a similar way we obtain the estimate from below

$$\int_\Gamma g(y, y/\varepsilon) d\sigma_y \geq \int_\Gamma \overline{g}_*(y, \nu_y) d\sigma_y - \eta_0.$$

As η_0 was arbitrary we have the two main estimates in the statement of the theorem.

Putting these together along with the IDDC we shall have the third statement. Indeed, due to IDDC we have that the set $\{x \in \Gamma : \nu_x \text{ rational}\}$ has zero surface measure, and the integral over this set is zero. For the rest of Γ we have irrational normals only, and hence the full averaging (Lemma 3.2) takes place and we obtain $\overline{g}_* = \overline{g}^*$. □

5. The case of Layered densities, almost periodicity, and ergodicity

In this section we shall deduce, similar results of that in Theorem 4.1 while replacing the periodicity assumption with layered materials/densities, almost-periodic and ergodic case. Nevertheless we shall only mention the results without deepening much into the analysis. The reader may easily verify the statements.

5.1. **Layered Materials.** If we assume the function $g(x,y)$ is independent of $(y_{k+1},\cdots,y_n)$ and is 1-periodic in $(y_1,\cdots,y_k)$ then one may naturally obtain results reminiscent of that of layered materials in homogenizations for PDE. Indeed one can obtain the following obvious result: *If the surface* Γ *does not have any flat parts in directions* e_i *(*$i=1,\cdots,k$*), then the averaging takes place.*

5.2. **Almost Periodic Case.** In the case of almost periodic functions (see [Bohr]) one obtains similar results by replacing the average integral $\overline{g}(x)$ with

$$\hat{g}(y)=\lim_{r\to\infty}\fint_{Q_r^+} g(x,y)dy.$$

The obvious details are left to the reader.

5.3. **Ergodic Case.** The results of our main theorem can be generalized further to the case of functions with ergodic properties (see [JKO]). Indeed, if we assume $g(x,y,\omega)$ is defined on $\Gamma\times\mathbb{R}^n\times D$, for some $D\subset\mathbb{R}^n$ and that $g(x,y,\omega)$ is statistically homogeneous field in (y,ω)-variable, i.e., $g(x,y,\omega)=h(x,T_y\omega)$ for some random variable $h(x,z)$ (random w.r.t. second variable) with underlying probability space and an ergodic n-dimensional dynamical system T_y. Hence $h(x,T_y\omega)$ admits an averaging

$$\lim_{\varepsilon\to0}\fint_K h(x,T_{y/\varepsilon}\omega)dx=\mathrm{Exp}(h(x,\cdot)).$$

One may now deduce similar results as that in Theorem 4.1 with

$$\overline{g}(z)=\mathrm{Exp}(g(z,\cdot,\cdot)).$$

6. Applications to Partial Differential Equations

6.1. **Dirichlet problem: Elliptic Case.** Let Ω be a bounded domain in $\mathbb{R}^n$, with piecewise smooth boundary. Let $g(x,y)$ be as before, and $f(x,y)$ have the same property as g. For simplicity, we shall assume

f, g are continuous in both variables (actually $L^2(\partial\Omega)$ would suffice). Let further $\Gamma_0 \subset \Omega$ be a piecewise C^1-curve, and define

$$\overline{\mu}^* = \overline{f}^* \chi_{\Gamma_0} d\sigma, \qquad \overline{\mu}_* = \overline{f}_* \chi_{\Gamma_0} d\sigma \qquad \overline{\mu} = \overline{f} \chi_{\Gamma_0} d\sigma.$$

Then the following result hold.

Theorem 6.1. *For a solution u_ε of (P_ε), we define (in Ω)*

$$u^*(x) := \limsup_{\varepsilon\to 0} u_\varepsilon(x) \qquad \text{and } u_*(x) := \liminf_{\varepsilon\to 0} u_\varepsilon(x)\ .$$

Then the following hold (in the weak sense):

$$\Delta\overline{u}^* \geq -\overline{\mu}^*, \qquad \Delta\overline{u}^* \leq -\overline{\mu}_* \qquad \text{in } \Omega,$$

$$u^*(x) \leq \overline{g}^*(x, \nu_x) \quad \text{on } \partial\Omega, \qquad u_*(x) \geq \overline{g}_*(x, \nu_x) \quad \text{on } \partial\Omega.$$

Moreover for any sequence of u_ε, there is a subsequence converging to a function u in Ω, satisfying

$$-\overline{\mu}^* \leq \Delta u \leq -\overline{\mu}_* \qquad \text{in the weak sense in } \Omega,$$

$$\overline{g}_*(x, \nu_x) \leq \varliminf_{z\to x} u(z) \leq \varlimsup_{z\to x} u(z) \leq \overline{g}^*(x, \nu_x), \qquad x \in \partial\Omega.$$

Proof. By Green's and Poisson's representations we have

$$u_\varepsilon(x) = \int_{\partial\Omega} P(x,y) g\left(y, \frac{y}{\varepsilon}\right) d\sigma_y + \int_{\Gamma_0} G(x,y) f\left(y, \frac{y}{\varepsilon}\right) d\sigma_y,$$

where P and G are Poisson respectively the Green functions of the domain.

By Theorem 4.1

$$u^*(x) \leq \int_{\partial\Omega} P(x,y)\overline{g}^*(y,\nu_y) d\sigma_y + \int_{\Gamma_0} G(x,y)\overline{f}^*(y,\nu_y) d\sigma_y,$$

and

$$u_*(x) \geq \int_{\partial\Omega} P(x,y)\overline{g}_*(y,\nu_y) d\sigma_y + \int_{\Gamma_0} G(x,y)\overline{f}_*(y,\nu_y) d\sigma_y,$$

which implies the first two statements in the theorem.

Now the last statement follows from the above inequalities, in an obvious way. ☐

To obtain an effective limit, in the above theorem, one needs to consider domains satisfying IDDC. In particular, using the above theorem and Lemma 3.2 (ii) we have the following result.

Corollary 6.2. *Let $\partial\Omega$ and Γ_0 satisfy IDDC, and u_ε be a solution to problem (P_ε). Then, u_ε converges to a function u in Ω, satisfying*

$$\Delta u = -\overline{\mu}(x), \quad \text{in } \Omega, \qquad u = \overline{g}(x) \quad \text{on } \partial\Omega.$$

The general nature of the method employed here suggests that we can apply this to situations where integral representations are possible. This can naturally go beyond the Dirichlet problem, or the Laplace operator, and as general as to systems, and equations of higher orders. We state explicitly that once one has an integral representation of any function, through kernel functions, then one may conclude similar statement as that of Theorem 6.1 and Corollary 6.2. We leave it to the reader to apply this to their favorite scenarios.

6.2. **Dirichlet problem: Parabolic Case.** Let us now consider the case of parabolic equations

$$\Delta u_\varepsilon - \partial_t u_\varepsilon = -\mu_\varepsilon(x,t) \quad \text{in } \Omega, \qquad u(x,t) = g_\varepsilon(x,t), \tag{9}$$

where $d\mu_\varepsilon(x,t) = f_\varepsilon(x,t)\chi_{\Gamma_0} d\sigma$, $f_\varepsilon(x,t) = f(x,t,x/\varepsilon,t/\varepsilon)$, $g_\varepsilon(x,t) = g(x,t,x/\varepsilon,t/\varepsilon)$, and $f(x,y,s,t)$, $g(x,y,s,t)$ are integrable in (x,s)-variables over Γ_0 and $\partial\Omega$ respectively, and 1-periodic in (y,t)-variable.

Here one may consider different cases, such as $\Omega = D\times(0,T)$, with D a domain in $\mathbb{R}^n$, or $\Omega \subset \mathbb{R}^n \times \mathbb{R}$ and time varying. Also Γ_0 can be take to be either time independent or varying in time.

Now a similar argument, using Poisson and Green representations, as in Section 6.1 can give us various type of results. In the case $\Omega = D \times (0,T)$ one obtains same type of results as that of Theorem 6.1. To obtain results of the nature of Corollary 6.2 one needs

- either to assume $\Omega = D \times (0,T)$, and D has IDDC condition along with f, and g being independent of their fourth variable, i.e. $f_\varepsilon(x,t) = f(x,x/\varepsilon,t)$, $g_\varepsilon(x,t) = g(x,x/\varepsilon,t)$
- or Ω and Γ_0, both have IDDC condition in $\mathbb{R}^{n+1}$.

In addition, Corollary 6.2 can be obtained similarly for $f_\varepsilon(x,t) = f(x,x/\varepsilon,t/\varepsilon^2)$, $g_\varepsilon(x,t) = g(x,x/\varepsilon,t/\varepsilon^2)$ with $\Omega = D \times (0,T)$, and D has IDDC condition.

6.3. **Neumann problem.** As mentioned at the end of the previous section, one can apply the technique of this paper to far reaching scenarios and problems, involving integral representations. The Neumann problem naturally fits into this category through Fredholm's alternative, and an integral representation. Indeed, let u_ε be a solution to the problem

$$\Delta u_\varepsilon = 0, \quad \text{in } \Omega, \qquad \partial_\nu u_\varepsilon = g(x, x/\varepsilon) \quad \text{on } \partial\Omega,$$

with g satisfying compatibility condition $\int_{\partial\Omega} g(y, y/\varepsilon) d\sigma_y = 0$ for all $\varepsilon > 0$. Then it is well-known that

$$u_\varepsilon(x) = \int_{\partial\Omega} F(x, y)\phi_\varepsilon(y) d\sigma_y,$$

where F is the Fundamental solution for Laplace operator (or the corresponding operator), and ϕ solves the Voltera integral equation of second kind, i.e.,

$$\phi_\varepsilon(x) = \int_{\partial\Omega} \partial_\nu F(x, y)\phi_\varepsilon(y) d\sigma_y + g(x, x/\varepsilon).$$

As ε tends to zero, ϕ_ε tends (weakly in $L^1(\partial\Omega)$) to a limit ϕ_0 solving

$$\phi_0(x) = \int_{\partial\Omega} \partial_\nu F(x, y)\phi_0(y) d\sigma_y + \overline{g}(x),$$

in a weak sense over the boundary of Ω. This happens exactly when the boundary of Ω has IDDC.

More accurately, the kernel of the bounded operator

$$T^\varepsilon(v) = v(x) - \int_{\partial\Omega} \partial_\nu F(x, y) v(y) d\sigma_y - g(x, x/\varepsilon),$$

acting on $L^1(\partial\Omega)$ space, is upper semi-continuous and has a unique element. In particular $\overline{\lim}_\varepsilon\, ker(T^\varepsilon) \subset ker(T^0)$, where

$$T^0(v) = v(x) - \int_{\partial\Omega} \partial_\nu F(x, y) v(y) d\sigma_y - \overline{g}(x).$$

By uniqueness of the solutions to the Fredholm operator this kernel must have only one element, and hence $\phi_\varepsilon \to \phi_0$, with $\phi_0 \in Ker(T^0)$. In other words ϕ_0 solves the Voltera equation above for the function $\overline{g}(x)$.

From here it follows that u_ε converges to $u_0 = \int_{\partial\Omega} F(x, y)\phi_0(y) d\sigma_y$, with ϕ_0 solving

$$\phi_0(x) = \int_{\partial\Omega} \partial_\nu F(x, y)\phi_0(y) d\sigma_y + \overline{g}(x).$$

Hence u_0 solves the averaged/effective Neumann problem

$$\Delta u_0 = 0, \quad \text{in } \Omega, \qquad \partial_\nu u_\varepsilon = \overline{g}(x) \quad \text{on } \partial\Omega.$$

A different way of analyzing this is to consider the solution of the Neumann problem in the weak form

$$\int_\Omega \nabla u_\varepsilon(y)\cdot\nabla\phi(y)dy = \int_{\partial\Omega} g(y, y/\varepsilon)\phi(y)d\sigma_y,$$

where ϕ is a test function in a reasonable class. Letting $\varepsilon \to 0$ we see that the integrals converge to

$$\int_\Omega \nabla u_0(y)\cdot\nabla\phi(y)dy = \int_{\partial\Omega} \overline{g}(y)\phi(y)d\sigma_y.$$

The latter in turn solves the Neuman problem with $\overline{g}(y)$ as the amount of flux at each boundary point.

7. Examples and illustrations

The behavior of the limit integrals in 2.1 are directly related to foliation of the fundamental cell Q_1^+. To illustrate this (in $\mathbb{R}^2$) consider a sequence $p_j = (-\sqrt{2}/j, 1)$. For $a \in [0, 1)$, let l_j^a be the line through the point $(0, a)$ and orthogonal to p_j. Then, due to the fact that p_j is rationally independent Lemma 3.2 implies

$$\lim_{\varepsilon\to 0} \fint_{l_j\cap Q_{\rho_\varepsilon}} g(x, y/\varepsilon)d\sigma_y = \overline{g}(x).$$

On the other hand $l_j \to l_0$ which is a line through $(0, a)$ and parallel to the x_1-axis. For the limit of the average for this line we then have

$$\lim_{\varepsilon\to 0} \fint_{l_0\cap Q_{\rho_\varepsilon}} g(x, y/\varepsilon)d\sigma_y = \overline{g}(x, \cdot, a).$$

where averaging takes place only on y_1-variable. In general $\overline{g}(x, \cdot, a) \neq \overline{g}(x)$. More importantly, we can in general not expect to take a very small ε_0 and expect

$$\fint_{l_0\cap Q_{\rho_\varepsilon}} g(x, y/\varepsilon_0)d\sigma_y \le \overline{g}^*(x),$$

for estimates from above, or $\ge \overline{g}_*(x)$, when considering estimates from below. The reader may easily verify this by taking $g(x, y) = g(y) = |\sin \pi y_1 \sin \pi y_2|$, and make computations to arrive at

$$\lim_{\varepsilon\to 0} \fint_{l_j\cap Q_{\rho_\varepsilon}} |\sin(\pi y_1/\varepsilon)\sin(\pi y_2/\varepsilon)|d\sigma_y = \frac{4}{\pi^2},$$

and

$$\lim_{\varepsilon\to 0} \fint_{l_0\cap Q_{\rho_\varepsilon}} |\sin(\pi y_1/\varepsilon)\sin(\pi y_2/\varepsilon)|d\sigma_y = \frac{2\sin(\pi a)}{\pi}.$$

Let us set

$$L_j^a = l_j^a \cap Q_{\rho_{\varepsilon_0}/\varepsilon_0}(mod\ 1).$$

Then one readily verifies that for ε_0 fixed, and very small

$$L_j^a \subset Q_1^+ \cap \left\{ x : a - \frac{M_{\varepsilon_0}\sqrt{2}}{j} \le x_2 \le a + \frac{M_{\varepsilon_0}\sqrt{2}}{j} \right\},$$

where $M_{\varepsilon_0} = \rho_{\varepsilon_0}/\varepsilon_0$. In particular, for j large L_j will never foliate the unit cell, and hence it is impossible to approximate the integral over $\partial\Omega$ by any covering, however small.

7.1. **Example 1.** We consider the case when g is periodic only in x_1-direction and when the domain is a slab with a unit normal direction ν.

For a $\nu \in \mathbb{S}^{n-1}$, set $\Omega = \{x : -R_1 < x \cdot \nu < R_2\}$. Let $g(x) = g(x_1)$ be independent of $(x_2, \cdots, x_n)$ and 1-periodic, i.e. $g(x+k) = g(x_1 + k_1) = g(x_1) = g(x)$ for $k \in \mathbb{Z}^n$. Now let u_ε be a solution of the following equation

$$\begin{cases} \triangle u_\varepsilon = 0 & \text{in } \Omega, \\ u_\varepsilon(x) = M & \text{on } x \cdot \nu = -R_1, \\ u_\varepsilon(x) = g\left(\frac{x}{\varepsilon}\right) & \text{on } x \cdot \nu = R_2. \end{cases} \tag{10}$$

We discuss three possible limits of u_ε whose homogenized equation can be found. Namely,

$$u^* = \limsup_{\varepsilon\to 0} u_\varepsilon, \quad u_* = \liminf_{\varepsilon\to 0} u_\varepsilon, \quad \text{and} \quad \overline{u} = \lim_{j\to\infty} u_{\varepsilon_i}, \tag{11}$$

where the subsequence u_{ε_i} is such that $E(u_{\varepsilon_i}) = \frac{1}{2}\int_\Omega |\nabla u_{\varepsilon_i}|^2 dx \to \liminf_{\varepsilon\to 0} E(u_\varepsilon)$ as $i \to \infty$ (See Figure 2). It turns out that u^*, u_*, and $\overline{u}$ don't follow simple homogenization whose boundary data is a simple average $\overline{g}$. It means there are nontrivial homogenization processes for each different limits. For $R \neq 0$, set

$$\overline{g}^*(R, e_1) = \max g, \quad \overline{g}_*(R, e_1) = \min g,$$

and

$$\overline{g}(R, M) = \begin{cases} \overline{g}^*(R, e_1) & \text{for } M > \sup g, \\ \overline{g}_*(R, e_1) & \text{for } M < \inf g, \\ M \quad \text{for } \inf g \le M \le \sup g. \end{cases} \tag{12}$$

If $R = 0$, let

$$\overline{g}^*(0, e_1) = \overline{g}_*(0, e_1) = \overline{g}(0, e_1) = g(0).$$

Proposition 7.1. *For the particular choice* $\nu = e_1$*, in equation (10), the limit functions* u^**,* u_**, and* $\overline{u}$ *will satisfy*

$$\begin{cases} \triangle u = 0 & \text{in } \Omega, \\ u(x) = M & \text{on } x \cdot e_1 = -R_1, \\ u(x) = A & \text{on } x \cdot e_1 = R_2, \end{cases} \tag{13}$$

where

$$A = \begin{cases} \overline{g}^*(R_2, e_1) & \text{if } u = u^*, \\ \overline{g}_*(R_2, e_1) & \text{if } u = u_*, \\ \overline{g}(R_2, e_1) & \text{if } u = \overline{u}. \end{cases} \tag{14}$$

Proof. Select ε_i such that $g\left(\frac{x_1}{\varepsilon_i}\right) = \max g$ and then $u_{\varepsilon_i} = u^*$ since all u_{ε_i} have the same boundary values. Hence it is clear that $\limsup_{\varepsilon \to 0} u_\varepsilon = u^*$. Similar argument can be applied to u_* and $\overline{u}$ to have the conclusion. □

Proposition 7.2. *In equation (10), when* $\nu \neq e_1$*,* u_ε *converges to* $\overline{u}$ *which is a solution to*

$$\begin{cases} \triangle u = 0 & \text{in } \Omega, \\ u(x) = M & \text{on } x \cdot \nu = -R_1, \\ u(x) = \overline{g} & \text{on } x \cdot \nu = R_2. \end{cases} \tag{15}$$

Proof. Choose any point x such that $x \cdot \nu = R_2$ and $\frac{x}{\varepsilon} = \tilde{x}$ (mod 1) for some $\tilde{x} \in Q_1$. Then, for $\nu \neq e_1$,

$$\begin{aligned} \fint_{\{(y-\tilde{x})\cdot\nu=0\}\cap Q_r} g\left(\frac{y}{\varepsilon}\right) d\sigma_y &= e_1 \cdot \nu \fint_{\{|y_1-\tilde{x}_1|\leq r\}} g\left(\frac{y_1}{\varepsilon}\right) \frac{1}{e_1 \cdot \nu} dy_1 \\ &= \fint_{\{|y_1-\tilde{x}_1|\leq r\}} g\left(\frac{y_1}{\varepsilon}\right) dy_1 \to \overline{g}, \qquad (\text{as } \epsilon \to 0), \end{aligned} \tag{16}$$

which is independent of the choice of $\tilde{x}$. It implies $\overline{g}^*(x,\nu) = \overline{g}_*(x,\nu) = \overline{g}(x)$. Let $P(x-y)$ be a Poisson Kernel of the the domain Ω. Then

$$u_\varepsilon(x) = M \int_{x\cdot\nu=-R_1} P(x-y)d\sigma_y + \int_{x\cdot\nu=R_2} P(x-y) g\left(\frac{y_1}{\varepsilon}\right) d\sigma_y.$$

Letting ϵ tend to zero and using first Lemma 3.3, and then Lemma 3.2 we shall have

$$\lim_{\epsilon \to 0} \int_{x\cdot\nu=R_2} P(x-y) g\left(\frac{y_1}{\varepsilon}\right) d\sigma_y = \int_{x\cdot\nu=R_2} P(x-y)\overline{g} d\sigma_y \, ,$$

which implies the conclusion. □



(a) fig 2.a (b) fig 2.b

Figure 1. *fig2.a shows the boundary* $x\cdot\nu = R_2$ *when* $\nu = e_1$ *or* $\nu \neq e_1$. *fig 2.b shows three possible limits (*u^*, u_* *and* u*) of* u_ε.

The next interesting question is to find the limit equation for the general converging sequence. In the following lemma, we will show there is a converging subsequence whose limit takes any value between the supremum of g and its infimum on $x \cdot e_1 = R_2$.

Proposition 7.3. *For any* A *such that* $\min g \leq A \leq \max g$, *there is a sequence* $\{u_{\varepsilon_i}\}$ *converging uniformly to* u *such that its limit* u *satisfies*

$$\begin{cases} \triangle u = 0 & \text{in } \Omega, \\ u(x) = M & \text{on } x \cdot e_1 = -R_1, \\ u(x) = A & \text{on } x \cdot e_1 = R_2. \end{cases} \tag{17}$$

Proof. There are $\varepsilon_i \to 0$ such that $g\left(\frac{x_1}{\varepsilon}\right) = A$ since g is 1-dimensional. Therefore all u_{ε_i} have the same boundary values, implying $u_{\varepsilon_i} = u(x)$. □

7.2. **Example 2.** In this example we confine ourselves to $\mathbb{R}^2$. We consider the case when g is periodic only in x_1-direction and the domain Ω is convex with two parallel flat parts of boundaries, orthogonal to $\nu = e_1$. For exactness we consider the following stadium like domain

$$\Omega = \left\{ x : |x_1| < R, |x_2| \leq 1 + \sqrt{R^2 - x_1^2} \right\}.$$

Let $g(x) = g(x_1)$ be 1-periodic, and independent of x_2-direction. Now let u_ε be a solution of the following equation

$$\begin{cases} \triangle u_\varepsilon = 0 & \text{in } \Omega, \\ u_\varepsilon(x) = g\left(\frac{x}{\varepsilon}\right) & \text{on } \partial\Omega, \end{cases} \tag{18}$$

then u^* and u_* (see (11)) are sub- and super-solutions of (18) respectively. In general, they are not solutions.

In the next result we state that the homogenized boundary data may not be continuous even though $g(x)$ is smooth.

Proposition 7.4. *There is a a smooth* 1*-periodic function* $g(x_1)$ *of one variable* x_1 *and a subsequence of solutions* $\{u_{\varepsilon_i}\}$ *to equation (18), converging to* u *such that* u *is not continuous on* $\partial\Omega$.

Proof. If $x_1 \neq \pm R$, then $x = (x_1, x_2) \in \partial\Omega$ satisfies IDD condition and then any limit will satisfy the boundary condition $u(x) = \overline{g}$. Now we select ε_i so that $g\left(\frac{x_1}{\varepsilon_i}\right) = \max g \neq \overline{g}$. Then u_{ε_i} has a converging subsequence to u which will be discontinuous at $x = (R, 1)$. □

REFERENCES

[AA] G. Allaire, and M. Amar; Boundary layer tails in periodic homogenization. ESAIM Control Optim. Calc. Var., 4:209243 (electronic), 1999.

[Bohr] H. Bohr, *Almost Periodic Functions*, Chelsea Publishing Company, New York, N.Y., 1947. ii+114 pp,

[BDLS] Barles, G., Da Lio, F.; Lions, P.-L.; Souganidis, P. E. Ergodic problems and periodic homogenization for fully nonlinear equations in half-space type domains with Neumann boundary conditions. Indiana Univ. Math. J. 57 (2008), no. 5, 23552375.

[CKL] S. Choi, I. Kim, K. Lee; Homogenization of Neumann boundary data with fully nonlinear operator (preprint).

[GM] Grard-Varet, David; Masmoudi, Nader; Homogenization and boundary layer, preprint.

[JKO] V. V. Jikov, S.M. Kozlov, O.A. Olenik, Homogenization of differential operators and integral functionals. (English summary) Translated from the Russian by G. A. Yosifian [G. A. Iosifyan]. Springer-Verlag, Berlin, 1994. xii+570 pp. ISBN: 3-540-54809-2.

SEOUL NATIONAL UNIVERSITY, SEOUL, 151-747, KOREA & KOREA INSTITUTE FOR ADVANCED STUDY, SEOUL,130-722, KOREA
E-mail address: `kiahm@snu.ac.kr`

DEPARTMENT OF MATHEMATICS, ROYAL INSTITUTE OF TECHNOLOGY, 100 44 STOCKHOLM, SWEDEN
E-mail address: `henriksh@kth.se`